## **Product Recommendation Guide**

| 1 - Low Risk   | 2 - Moderate Risk  | 3 - High Risk  |
|--|--|--|
| Gel CTX4<br>Gel S000   | TX3 Rinse  TX3  Rinse  TX3  TX3  TX3  TX3  TX3  TX3  TX3  TX   | TR Treatment Rinse   |
| CTx12 Kit or CTx4 Gel 1100 (optional)  | CTx21 Kit  | CTx36 Kit  |
| Brush CTx4 Gel 5000 or CTx4 Gel 1100  Replaces current toothpaste Brush twice daily with a pea-sized amount until product is gone  Avoid eating or drinking for 30 minutes after use | Rinse 1st CTx3 Rinse  • Use twice daily until product is gone  • Swish for 60 seconds with 10 mL  Brush 2nd CTx4 Gel 5000  • Replaces current toothpaste  • Brush twice daily with a pea-sized amount until product is gone  • Avoid eating or drinking for 30 minutes after use | Rinse 1st CTx4 Treatment Rinse  Use twice daily until product is gone  Mix 5 mL of Solution A with 5 mL of Solution B in the provided mixing cup  Swish for 60 seconds  Brush 2nd CTx4 Gel 5000  Replaces current toothpaste  Brush twice daily with a pea-sized amount until product is gone  Avoid eating or drinking for 30 minutes after use |